Nutrient Broth

Liquid medium for the cultivation of nonfastidious microorganisms.

DESCRIPTION

Nutrient Broth is a liquid medium used for the cultivation of a wide variety of organisms from clinical specimens and other materials.

This medium can be enriched with other ingredients such as blood, serum, sugars, etc., for special purposes.

| TYPICAL FORMULA | (g/ l) |
|----------------------------|----------------|
| Beef Extract | 1.0 |
| Peptone | 5.0 |
| Yeast Extract | 2.0 |
| Sodium Chloride | 5.0 |
| Final pH 6.8 ± 0.2 at 25°C | |

METHOD PRINCIPLE

Beef extract and peptone provide amino acids, nitrogen, carbon, vitamins and minerals for organisms growth. Yeast extract is a source of vitamins, particularly of B-group. Sodium chloride maintains the osmotic balance of the medium.

PREPARATION

Dehydrated medium

Suspend 13 g of the powder in 1 liter of distilled or deionized water. Mix well. Heat to boil shaking frequently until completely dissolved. Sterilize in autoclave at 121°C for 15 minutes.

TEST PROCEDURE

Inoculate broth with test sample. Incubate at $35 \pm 2^{\circ}$ C for 18-24 hours or longer if necessary.

INTERPRETING RESULTS

Turbidity indicates microbial growth.

APPEARANCE

Dehydrated medium: free-flowing, homogeneous, white to light beige.

Prepared medium: clear to slightly opalescent, light amber.

STORAGE

The powder is very hygroscopic, store the powder at 10-30°C, in a dry environment, in its original container tightly closed. Store bottles and tubes at 10-25°C away from light. Do not use the product beyond its expiry date on the label or if product shows any evidence of contamination or any sign of deterioration.

SHELF LIFE

Dehydrated medium: 4 years. Medium in tubes/bottles: 2 years.

Distribué par :

Z.A de Gesvrine - 4 rue Képler - B.P.4125 44241 La Chapelle-sur-Erdre Cedex - France t.:+33 (0)2 40 93 53 53 | f.:+33 (0)2 40 93 41 00 commercial@humeau.com



QUALITY CONTROL

The medium is inoculated with the microbial strains indicated in the QC table.

Inoculum for productivity: ≤100 CFU

Incubation conditions: aerobically at $35 \pm 2^{\circ}$ C for 18-24 hours.

QC Table.

| Microorganism | | Growth |
|-----------------------|-------------|--------|
| Escherichia coli | ATCC® 25922 | Good |
| Staphylococcus aureus | ATCC® 25923 | Good |

WARNING AND PRECAUTIONS

The product does not contain hazardous substances in concentrations exceeding the limits set by current legislation and therefore is not classified as dangerous. It is nevertheless recommended to consult the safety data sheet for its correct use. The product is intended for *In vitro* diagnostic use and must be used only by properly trained operators.

DISPOSAL OF WASTE

Disposal of waste must be carried out according to national and local regulations in force.

BIBLIOGRAPHY

- Association of Official Analytical Chemists (1995) Official methods of analysis of AOAC International, 16th ed.
- 2. Marshall, R.T. (ed.) (1993) Standard methods for the microbiological examination of dairy products, 16th ed.
- 3. American Public Health Association (1923) Standard methods of water analysis, 5th ed.

| PRESENTATION | | Contents | Ref. |
|----------------|-------------------|--------------------|---------|
| Nutrient Broth | Tubes | 20 x 10 ml tubes | 24103 |
| Nutrient Broth | Tubes | 50 x 5 ml tubes | 27503 |
| Nutrient Broth | Bottles | 6 x 100 ml bottles | 402000 |
| Nutrient Broth | Bottles | 6 x 500 ml bottles | 470050 |
| Nutrient Broth | Dehydrated medium | 500 g of powder | 610037 |
| Nutrient Broth | Dehydrated medium | 100 g of powder | 620037 |
| Nutrient Broth | Dehydrated medium | 5 kg of powder | 6100375 |

TABLE OF SYMBOLS

| LOT Batch code | IVD In vitro Diagnostic Medical Device | Manufacturer | S Use by | Fragile, handle with care |
|----------------------|---|---------------------------------------|---|---------------------------|
| REF Catalogue number | Temperature limitation | Contains sufficient for <n> tests</n> | Caution, consult Instruction For Use | Do not reuse |

