

Technical Data Sheet

WL Nutrient Agar

Ordering number: 1.10866.0500

For the enumeration and cultivation of yeasts and bacteria in the microbiological control carried out in the brewing and other fermentation industries (GREEN and GRAY 1950, 1951; GRAY 1951).

GRAY (1951) has given a detailed description of the use of WL Nutrient Agar and WL Differential Agar in the microbiological quality control employed in the beer-brewing industry.

Mode of Action

WL (Wallerstein Laboratories) Nutrient Agar has a pH of 5.5, which is optimal for the enumeration of brewers' yeast. If bakers' or distillers' yeast is to be examined, the pH should be adjusted to 6.5 (better yields). When cultivating microorganisms from an alcoholic mash, tomato juice should be added to the medium. WL differential agar contains cycloheximide to suppress yeasts and any moulds which may be present; this medium allows reliable counting of all bacteria which may be encountered in the tests performed in brewery laboratories.

Typical Composition (g/L)

WL Nutrient Agar	
Yeast extract	4.0
Casein hydrolysate	5.0
D(+)-glucose	50.0
Potassium dihydrogen phosphate	0.55
Potassium chloride	0.425
Calcium chloride	0.125
Magnesium sulfate	0.125
Iron(III) chloride	0.0025
Manganese sulfate	0.0025
Bromocresol green	0.022
Agar-agar**	17.0

**Agar-agar is equivalent to other different terms of agar.

Preparation

Suspend 77 g/litre, if required dissolve the medium in a mixture of 400 ml clarified tomato juice and 600 ml demineralized water, adjust the pH to 6.5 if necessary, autoclave (15 min at 121 °C), pour plates.

pH: 5.5 ± 0.2 at 25 °C.

The plates are clear and blue-green.

Experimental Procedure and Evaluation

Dilute the sample material, spread 0.1 ml on WL Nutrient Agar and, if necessary, on WL differential agar.

Incubation: up to 2 weeks at 25 ° and, if applicable, at 30 °C aerobically. WL Differential Agar should be incubated both aerobically and anaerobically.

Count the number of colonies per plate and calculate the microbial count. Acetic acid bacteria, flavobacteria, thermobacteria, Proteus bacteria and other species grow on WL Differential Agar under aerobic conditions whereas lactobacilli and pediococci proliferate under anaerobic conditions.

Storage

Store at +15 °C to +25 °C, dry and tightly closed. Do not use clumped or discolored medium. Protect from UV light (including sun light). For *in vitro* use only.

Quality Control

Control strains	Growth, 4 days, 28°C; aerobic
Candida albicans ATCC 10231 (WDCM 00054)	Good to very good
Saccharomyces cerevisiae ATCC 9763 (WDCM 00058)	Good to very good
Lactobacillus acidophilus ATCC 4356 (WDCM 00098)	Fair to very good
Leuconostoc mesenteroides ATCC 9135 (WDCM 00108)	Good to very good
Enterococcus faecalis ATCC 11700	Good to very good
Escherichia coli ATCC 11775 (WDCM 00090)	Good to very good
Proteus mirabilis ATCC 29906 (WDCM 00023)	Good to very good

Please refer to the actual batch related Certificate of Analysis.



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

Merck Millipore, the M logo, Anaeroclip, Anaerocult, Anaerotest are registered trademarks of Merck KGaA, Darmstadt, Germany. ATCC is a registered trademark of ATCC, Manassas, VA, USA. Lit. No. TN1823EN00

Literature

GRAY, P.P.: Some advances in microbiological control for beer quality. - **Wallerstein Lab. Comm.**, **14**; 169-183 (1951).

GREEN, S.R., a. GRAY, P.P.: Paper read at Am. Soc. of Brewing Chemists Meeting; - **Wallerstein Lab. Comm.**, **12**; 43 (1950).

GREEN, S.R., a. GRAY, P.P.: A differential procedure applicable to bacteriological investigation in brewing. - **Wallerstein Lab. Comm.**, **13**; 357-366 (1950).

GREEN, S.R., a. GRAY, P.P.: A differential procedure for bacteriological studies useful in the fermentation industries. - **Wallerstein Lab. Comm.**, **14**; 289-295 (1951).

Ordering Information

Product	Cat. No.	Pack size
WL Nutrient Agar	1.10866.0500	500 g
Anaeroclip®	1.14226.0001	1 x 25
Anaerocult® A mini	1.01611.0001	1 x 25
Anaerocult® P	1.13807.0001	1 x 25
Anaerotest®	1.15112.0001	1 x 50

Merck KGaA, 64271 Darmstadt, Germany
Fax: +49 (0) 61 51 / 72-60 80
mibio@merckgroup.com
www.merckmillipore.com/biomonitoring

Find contact information for your country at:
www.merckmillipore.com/offices
For Technical Service, please visit:
www.merckmillipore.com/techservice



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

Merck Millipore, the M logo, Anaeroclip, Anaerocult, Anaerotest are registered trademarks of Merck KGaA, Darmstadt, Germany. ATCC is a registered trademark of ATCC, Manassas, VA, USA. Lit. No. TN1823EN00